

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 May 2001 (17.05.2001)

PCT

(10) International Publication Number
WO 01/34561 A1

- (51) International Patent Classification⁷: C07C 303/40, 311/35, C07D 209/14 (74) Agent: GOLDSCHIED, Bettina; c/o BASF Aktiengesellschaft, 67056 Ludwigshafen (DE).
- (21) International Application Number: PCT/EP00/10581 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 27 October 2000 (27.10.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
9926250.3 6 November 1999 (06.11.1999) GB
- (71) Applicant (*for all designated States except US*): KNOLL AKTIENGESELLSCHAFT [DE/DE]; 67061 Ludwigshafen (DE).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): HOLMAN, Nicholas, John [GB/GB]; R5 Pennyfoot Street, Nottingham NG1 1GF (GB). FRIEND, Christopher, Lyndon [GB/GB]; R4 Pennyfoot Street, Nottingham NG1 1GF (GB).

Published:


— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.


(54) Title: PROCESSES FOR THE PREPARATION OF SUMATRIPTAN AND RELATED COMPOUNDS




(A)




(B)



(C)



(I)



X⁻ (II)

(57) Abstract: The present invention relates to a process for preparing phenylhydrazines of formula (I) in which R represents CH₂SO₂NHCH₃, CH₂CH₂SO₂Ph, CH₂CH₂SO₂NHMe or a group of structure (A), (B), (C), in which a diazonium salt of formula (II) in which X represents an anion derived from hydrochloric acid, sulphuric acid, acetic acid, phosphoric acid, tetrafluoroboric acid or hydrobromic acid is reduced by a dithionite salt. The resulting phenylhydrazines can be converted to the corresponding indole derivatives by the Fischer indole synthesis.

The present invention relates to processes for preparing phenylhydrazines which are useful intermediates in the preparation of indoles which are useful as therapeutic agents.


Several indole derivatives are currently on the market or under development as pharmaceuticals. For example, sumatriptan (Glaxo Wellcome) 3-[2-(dimethylamino)ethyl]-*N*-methyl-1*H*-indole-5-methanesulphonamide is currently used in the treatment of migraine. This compound is described in US 4,816,470 and US 5,737,845.

Other indoles of pharmaceutical interest are: almotriptan (WHO Drug Information, Vol.10, No. 4, [1996]), avitriptan (WHO Drug Information, Vol.10, No. 4, [1996]), eletriptan (WHO Drug Information, Vol.9, No. 4, [1995]), frovatriptan (Br.J. Pharmacol. [119, Proc. Suppl., 109P, 1996]), naratriptan, rizatriptan (WHO Drug Information, Vol.10, No. 2, [1996]) and zolmitriptan (J. Med. Chem. [38, No.18, 3566-80, 1995]).


In the preparation of such indoles phenylhydrazines are key intermediates which may be cyclised into indoles using the well known Fischer indole synthesis. In US 4,816,470 the method of preparation of such compounds involves the diazotisation of an aniline followed by reduction of the diazonium salt obtained with stannous chloride dihydrate. We have found that the use of tin reagents in this reduction presents a number of problems. Firstly there are environmental issues relating to the disposal of toxic wastes containing tin residues. Secondly, low temperature vessels are required to carry out the reduction and thirdly it is often difficult to remove the last traces of tin salts from the intermediate and from later stages of the reaction sequence.

Surprisingly a process has been found which uses a cheaper reducing agent which causes minimal environmental problems. In addition the process does not require the use of low temperature vessels, allows the telescoping of the process, i.e. that is the combination of more than one step, and increases the purity of the intermediate obtained and further products in the reaction sequence.


The present invention provides a process for preparing a compound of formula I including salts thereof



wherein R represents $\text{CH}_2\text{SO}_2\text{NHCH}_3$, $\text{CH}_2\text{CH}_2\text{SO}_2\text{Ph}$, $\text{CH}_2\text{CH}_2\text{SO}_2\text{NHMe}$ or a group of structure:




comprising the reduction of a diazonium salt of formula II



in which R is as previously defined and X^- represents an anion derived from hydrochloric acid, sulphuric acid, acetic acid, phosphoric acid, tetrafluoroboric acid or hydrobromic acid, with a dithionite salt.


In another aspect the present invention provides a process for preparing a compound of formula I or a salt thereof




in which R is as previously defined comprising the steps of

3

- a) reacting a compound of formula III or salt thereof



in which R is as previously defined with a diazotising agent optionally in the presence of acid to give a diazonium salt of formula II




in which R and X⁻ are as previously defined and

- b) reducing the diazonium salt with a dithionite salt to give the compound of formula I.

Suitable salts of the compounds of formula I and III include acid addition salts formed with organic or inorganic acids for example hydrochlorides, hydrobromides, sulphates, nitrates, phosphates, formates, mesylates, citrates, benzoates, fumarates, maleates, toluene-p-sulphonates and succinates. Preferably the salt is the hydrochloride or hydrobromide salt.


In a further aspect the present invention provides a process in which the compound of formula III or salt thereof as previously defined is prepared by reacting a compound of formula IV




in which R is as previously defined with a reducing agent, optionally in the presence of an acid or with an additional salt formation step eg reaction with an acid, where a salt of the compound of formula III is required. Preferably the compound of formula III is used without isolation.

Consequently the present invention provides a process for the preparation of a compound of formula I which comprises the steps of:

- a) reacting a compound of formula IV




in which R is as previously defined with a reducing agent to give a compound of formula III




in which R is as previously defined and

- b) reacting the compound of formula III with a diazotising agent optionally in the presence of acid to give a diazonium salt of formula II




in which R and X⁻ are as previously defined and


- c) reducing the diazonium salt with a dithionite salt to give the compound of formula I.




In a further aspect the present invention comprises a process for preparing a compound of formula V



including pharmaceutically acceptable salts thereof in which R is as previously defined and R1 is a group of formula a), b), c), d), or e)




comprising performing a Fischer Indole synthesis by reacting a compound of formula I with an aldehyde of formula VI in which R1 is as previously defined optionally in the presence of an acid catalyst.



VI


Optionally a protected form of the aldehyde such as an acetal may be used this reaction. Optionally an aldehyde or a protected form of the aldehyde of formula VII may be used where R2 is a group capable of being transformed into a group R1 upon cyclisation to a compound of formula V, for instance when R2 is $-\text{CH}_2\text{CH}_2\text{Cl}$ to give compounds of formula V with R1 is $-\text{CH}_2\text{CH}_2\text{NH}_2$ using the well known Grandberg version of the Fischer Indole synthesis.



VII

Optionally it may be preferred to synthesise one compound of formula V by reacting another compound of formula V, for instance, when R1 is $\text{CH}_2\text{CH}_2\text{NMe}_2$ by reductive alkylation of the compound V where R1 is $\text{CH}_2\text{CH}_2\text{NH}_2$

Optionally it may be preferred to isolate the intermediate hydrazone VIII



VIII

In which R and R1 are as previously defined prior to performing the cyclisation to V.

Suitably the diazotising agent is a metal nitrite salt or alkyl nitrite. Preferably the diazotising agent is sodium nitrite or butyl nitrite. Most preferably the diazotising agent is sodium nitrite.

Suitably the diazotising agent is present in the range of 0.5–3 molar equivalents with respect to the compound of formula II. Preferably the diazotising agent is present in the range of 0.8–1.5 molar equivalents with respect to the compound of formula II. More preferably the diazotising agent is present in the range of 0.9–1.1 molar equivalents with respect to the compound of formula II.

Optionally it may be preferred to add a reagent, for example sulphamic acid, to destroy excess nitrous acid at the end of this step.

Suitably the acid is hydrochloric acid, sulphuric acid, acetic acid, phosphoric acid, tetrafluoroboric acid or hydrobromic acid. Preferably the acid is hydrochloric acid.

Suitably the acid is present in the range of 1–10 molar equivalents with respect to the compound of formula II. Preferably the acid is present in the range of 2–8 molar equivalents with respect to the compound of formula II. More preferably the acid is present in the range of 3–6 molar equivalents with respect to the compound of formula II.

Suitably the dithionite salt is a metal dithionite salt. Most preferably the dithionite salt is sodium dithionite.

Suitably the dithionite is present in the range of 1–5 molar equivalents with respect to the compound of formula II and is used as an aqueous solution or suspension in the presence of base, preferably sodium hydroxide. Preferably the dithionite is present in the range of 2–4 molar equivalents with respect to the compound of formula II and is used as an aqueous solution or suspension in the presence of base, preferably sodium hydroxide. Preferably isopropanol can be used as a co-solvent to reduce foaming.


Suitably the diazotisation step is carried out at a temperature in the range of -20°C to $+20^{\circ}\text{C}$. Preferably the diazotisation step is carried out at a temperature in the range of -10°C to $+10^{\circ}\text{C}$. More preferably the diazotisation step is carried out at a temperature in the range of -5°C to $+5^{\circ}\text{C}$.

Suitably the reduction of the diazonium salt is carried out at a temperature in the range of -50°C to $+50^{\circ}\text{C}$. Preferably the reduction of the diazonium salt is carried out at a temperature in the range of -10°C to $+30^{\circ}\text{C}$. More preferably the reduction of the diazonium salt is carried out at a temperature in the range of -5°C to $+25^{\circ}\text{C}$.

Suitably the reducing agent for the nitro compound is hydrogen in the presence of a catalyst, sodium dithionite, iron in the presence of acid or lithium aluminium hydride. Preferably the reducing agent for the nitro compound is hydrogen in the presence of a catalyst, sodium dithionite, or lithium aluminium hydride. Most preferably the reducing agent for the nitro compound is hydrogen in the presence of a palladium catalyst.

⇒ The hydrogenation/diazonium reaction/reduction can be carried out as a one-pot reaction.

In a further aspect the present invention provides a process in which a compound of formula V




in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and R1 is $\text{CH}_2\text{CH}_2\text{NMe}_2$, is prepared by reacting a compound

NO Si


of formula V, in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and R1 is $\text{CH}_2\text{CH}_2\text{NH}_2$, with a reducing agent and a formaldehyde equivalent in the presence of a buffer.

In another aspect the present invention provides a process for the preparation of sumatriptan or a pharmaceutically acceptable salt thereof comprising the following steps:

a) reducing a compound of formula




in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ with a reducing agent, optionally in the presence of an acid to give a compound of formula III



or optionally a salt thereof in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and


b) reacting the compound of formula III with a diazotising agent optionally in the presence of acid to give a compound of formula II




in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and X^- represents a chloride, bromide, acetate, hydro-sulphate or phosphate anion and

c) reducing the compound of formula II with a dithionite salt to give a compound of formula I

in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and




- d) reacting the compound of formula I in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ with an aldehyde of formula VII



VII

in which R_2 represents $\text{ClCH}_2\text{CH}_2-$ and the aldehyde group is protected as an acetal (preferably the dimethyl acetal) in the presence of a buffer optionally in the presence of an acid catalyst to give a compound of formula V



or a salt thereof in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and R_1 is $\text{CH}_2\text{CH}_2\text{NH}_2$ and

- e) reacting the compound of formula V obtained in d) with a formaldehyde equivalent and a reducing agent in the presence of a buffer to give a compound of formula V in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and R_1 is $\text{CH}_2\text{CH}_2\text{NMe}_2$ or a pharmaceutically acceptable salt thereof.

Preferably none of the intermediate compounds is isolated in this process.

Suitably the reducing agent is a hydride equivalent such as sodium borohydride, sodium cyanoborohydride, sodium triacetoxyborohydride and lithium aluminium hydride. Preferably the reducing agent is sodium borohydride or sodium cyanoborohydride. Most preferably the reducing agent is sodium borohydride.

Suitably the reducing agent is present in the range of 0.25 – 5 molar equivalents with respect to the compound of formula V. Preferably the reducing agent is present in the range of 0.5 – 4 molar equivalents with respect to the compound of formula V. More preferably the

reducing agent is present in the range of 0.75 – 3 molar equivalents with respect to the compound of formula V.

Suitably the formaldehyde equivalent is formaldehyde, paraformaldehyde or dimethoxymethane. Preferably the formaldehyde equivalent is formaldehyde or paraformaldehyde. Most preferably the formaldehyde equivalent is formaldehyde as an aqueous solution.

Suitably the formaldehyde equivalent is present in the range of 0.5 – 18 molar equivalents with respect to the compound of formula V. Preferably the formaldehyde equivalent is present in the range of 1 – 10 molar equivalents with respect to the compound of formula V. More preferably the formaldehyde equivalent is present in the range of 1.9 – 5 molar equivalents with respect to the compound of formula V.

Suitably the buffer used keeps the pH of the reaction solution between pH 6 and pH 14. Preferably the buffer keeps the pH of the reaction solution between pH 7 and pH 11. More preferably the buffer keeps the pH of the reaction solution between pH 8 and pH 10. Most preferably the buffer is sodium hydrogenphosphate.

Suitably the buffer is present in the range of 0.1 – 10 molar equivalents with respect to the compound of formula V. Preferably the buffer is present in the range of 0.2 – 5 molar equivalents with respect to the compound of formula V. Most preferably the buffer is present in the range of 0.5 – 3 molar equivalents with respect to the compound of formula V.

Suitable pharmaceutically acceptable salts of the compound of formula V (including sumatriptan) include acid addition salts formed with organic or inorganic acids for example hydrochlorides, hydrobromides, sulphates, nitrates, phosphates, formates, mesylates, citrates, benzoates, fumarates, maleates and succinates. Other salts may be useful in the preparation of the compound of formula I e.g. creatinine sulphate adducts, and salts with e.g. toluene-p-sulphonic acid. When the compound of formula V is sumatriptan the salt is preferably the succinate salt or the hemisulphate salt.

The invention is illustrated by the following Examples which are given by way of example only. The final products of each of these Examples were characterised by one or more of the following procedures: high performance liquid chromatography, elemental analysis, nuclear magnetic resonance spectroscopy, mass spectroscopy and infrared spectroscopy.

11
EXAMPLES

Example 1

- a) A stirred mixture of 4-amino-*N*-methylbenzenemethanesulphonamide (35.6 g), concentrated hydrochloric acid (83.7 ml) and water (314 ml) was heated at 50°C for 15 minutes and the solution then cooled to -5°C. A solution of sodium nitrite (12.5 g) in water (21 ml) was then added dropwise over 10 minutes. The resulting solution was stirred for 1 hour and then added over 10 minutes to a stirred suspension of sodium dithionite (sodium hydrosulphite) (96.1 g) in water (420 ml) and 46/48% w/w sodium hydroxide solution (34 ml) at -5°C to +5°C. The suspension was stirred for 2.75 hours. 46/48% w/w Sodium hydroxide (22 ml) was added and the mixture was stirred for 20 minutes at 20°C and then for 40 minutes at 0-5°C. The mixture was filtered and the product was washed with water (3 x 110 ml) and dried to give the free base of the product.

The free base of the product (31.5 g) and ethanol (315 ml) were stirred and heated to reflux. Concentrated hydrochloric acid (12.3 ml) was added followed by ethanol (155 ml). The mixture was boiled under reflux for 10 minutes and then cooled to 0°C. The product was collected by filtration, washed with ethanol (155 ml) and dried to give 4-hydrazino-*N*-methylbenzenemethanesulphonamide hydrochloride, yield 31.4 g.

- b) Absolute ethanol (50 ml) was added to a stirred suspension of 4-hydrazino-*N*-methylbenzenemethanesulphonamide hydrochloride (6.42 g) in water (20ml) and the mixture was stirred for 10 minutes. 4-Chlorobutanal dimethyl acetal (3.85 g) was added and washed in with more absolute ethanol (11.7 ml). 2M Hydrochloric acid (0.22 ml) was added and the solution was stirred at ambient temperature for 4.5 hours. Sodium hydrogenphosphate (3.01 g) was added and the mixture was stirred at ambient temperature for 10 minutes and then gradually heated to boiling under reflux over 40 minutes. The mixture was stirred and boiled under reflux for a further 3 hours and then allowed to stand for 16 hours at ambient temperature. The mixture was concentrated under reduced pressure (around 60 ml removed) and then dichloromethane (25 ml) and water (25 ml) were added, followed by potassium carbonate (0.74 g). At this point the pH was 5. The mixture was filtered and the filtrate was separated. The aqueous layer was washed with more dichloromethane

(2 x 25 ml) and the combined organic layer and washings were evaporated to dryness under reduced pressure to give a solid by-product, yield 1.60 g.

The aqueous layer was mixed with dichloromethane (125 ml), absolute ethanol (60 ml) potassium carbonate (37 g) and water (12 ml). The mixture was stirred for 35 minutes and then separated. The organic layer was treated with charcoal and stirred at ambient temperature for 1 hour. The mixture was filtered and the filtrate was concentrated to a mass of 20 g under reduced pressure. The mixture was stirred for 1 hour and isopropyl acetate (62 ml) was added and the suspension was stirred for 64 hours. The solid was collected by filtration, washed with more isopropyl acetate (10 ml) and dried under vacuum at 50°C to give 3-(2-aminoethyl-*N*-methyl-1*H*-indole-5-methanesulphonamide, yield 3.64 g (53.4%). Purity 93.63%.

Comparative Example

4-Hydrazino-*N*-methylbenzenemethanesulphonamide hydrochloride, which had been prepared by diazotisation of 4-amino-*N*-methylbenzenemethanesulphonamide with sodium nitrite and then reduced with tin chloride, was reacted according to the procedure of Example 2. 3-(2-Aminoethyl-*N*-methyl-1*H*-indole-5-methanesulphonamide was obtained in a yield of 48.8% and was found to be 92.25% pure by HPLC.

Example 2

N-Methyl-4-nitrobenzenemethanesulphonamide (46.0 g, 0.23 mol), 10% palladium on carbon (4.6 g) and 2M hydrochloric acid (180 ml) in water (200 ml) were stirred for 1.5 hours under 2.5 atmospheres of hydrogen at 20°C. The reaction mixture was then filtered through a celite bed and washed with further water (100 ml). A portion of this filtrate (50 ml) was taken and then diluted with concentrated hydrochloric acid (12.7 ml). The stirred suspension was cooled to below 0°C and a solution of sodium nitrite (2.2 g, 32 mmol) in water (4 ml) added dropwise over 20 minutes under an atmosphere of nitrogen. After stirring for 15 minutes, the clear solution was transferred *via* a cannula to a solution of sodium dithionite (17.0 g, 98 mmol) and 46/48% w/w sodium hydroxide (5.9 ml) in water (75 ml) at -5°C. The mixture was then warmed to room temperature and stirred for a further 2.5 hours. 46/48% w/w sodium hydroxide (6.5 ml) was then added until the pH of the solution was approximately 9. Stirring at room temperature was continued for a further 0.5 hours

followed by cooling in an ice bath for 0.5 hours. Filtration of the reaction mixture gave 4-hydrazino-*N*-methylbenzenemethanesulphonamide as an off-white solid (2.5 g, 48%)

Example 3

A stirred mixture of 4-amino-*N*-methylbenzenemethanesulphonamide (50.0 g, 0.25 mmol), conc. hydrochloric acid (117 ml) and water (356 ml) was heated at 50°C for 15 minutes and the solution then cooled to -5°C. A solution of sodium nitrite (17.5 g) in water (30 ml) was then added dropwise over 15 minutes and the resulting solution stirred for a further 15 minutes. The reaction mixture was then added via a cannula to a stirred suspension of sodium dithionite (135 g, 0.66 mol), water (365 ml), 46/48% w/w sodium hydroxide solution (23.5 ml) and IPA (40 ml) at -5°C. The temperature was kept around -5°C during the 40 minute addition. The suspension was then warmed to room temperature and stirred for 2.5 hours before 46/48% w/w Sodium hydroxide (53.5 ml) was added to give a pH of 7-8. Finally, the mixture was stirred for 30 minutes, product filtered and then washed with water (123 ml) to give a cream solid.

To form the hydrochloride salt, conc. hydrochloric acid (20 ml) was added to the free base suspended in isopropanol (400 ml) at ambient. After stirring for 15 minutes, the product was collected by filtration, washed with isopropanol (125 ml) and dried to give 4-hydrazino-*N*-methylbenzenemethanesulphonamide hydrochloride (45.7 g, 73% at 97% purity by HPLC).

Example 4

N-Methyl-4-nitrobenzenemethanesulphonamide (23.0 g, 0.1 mol), 5% palladium on carbon (9.4 g) and conc. hydrochloric acid (11 ml) in water (245 ml) were stirred for 2 hours under 5 atmospheres of hydrogen at 50°C. The reaction mixture was then filtered through a celite bed and washed with further conc. hydrochloric acid (6 ml) in water (34 ml). The filtrate was then diluted with conc. hydrochloric acid (30 ml) and the solution cooled to -5°C to give a suspension. Sodium nitrite (7.0 g, 0.101 mol) in water (12 ml) was then added dropwise over 20 minutes keeping the temperature around -5°C. After stirring for 15 minutes, the clear solution was transferred via a cannula to a solution of sodium dithionite (54.0 g, 0.264 mol), 46/48% w/w sodium hydroxide (9.4 ml), water (197 ml) and isopropanol (20 ml) at -5°C. The temperature was kept around -5°C during the 40 minutes addition. The mixture was then warmed to room temperature and stirred for a further 2.5 hours. 46/48% w/w sodium hydroxide (21.4 ml) was then added until the pH of the solution was approximately

7.8. Stirring at room temperature was continued for a further 0.5 hours followed by filtration of the reaction mixture, washing with water (50 ml) and then isopropanol (150 ml) to give 4-hydrazino-N-methylbenzenemethanesulphonamide as an off-white solid. Forming the hydrochloride salt as above, gave an overall yield of 77% at 97.6% purity by HPLC.

Example 5

3-(2-Aminoethyl)-N-methyl-1H-indole-5-methanesulphonamide (5.0 g, 18.7 mmol), prepared by the method of Example 1, and sodium hydrogenphosphate (5.0 g, 35.2 mmol) were heated to 40°C in methanol (50 ml) for 15 minutes and then recooled to room temperature. Solutions of 37% aqueous formaldehyde (5 ml) and sodium borohydride (0.72 g) in water (5 ml stabilised with one drop of 46/48% w/w sodium hydroxide) were added simultaneously over one hour at a temperature between 17 and 21°C. The mixture was stirred at room temperature for a further 0.5 hours, then filtered and the filter bed washed with methanol (10 ml). The combined filtrates were then adjusted to pH 6 by addition of 4M hydrochloric acid, concentrated under reduced pressure (to approximately 20 g) and acidified to pH 1 with more 4M hydrochloric acid. Ethyl acetate (30 ml) was added and then potassium carbonate was added to give a pH about 11 and the product extracted in a separating funnel. The aqueous layer was further extracted with ethyl acetate (30 ml) and the combined organic layers dried over sodium sulphate, filtered and concentrated to give sumatriptan free base (4.7 g, 85%, HPLC showed 87% compound).

This material was of suitable quality for conversion into sumatriptan mono-succinate or sumatriptan hemisulphate as described in GB 2,162,522 and EP 490,689 respectively.

Example 6


3-(2-Aminoethyl)-N-methyl-1H-indole-5-methanesulphonamide hydrochloride (50.0 g, 0.165 mol) and sodium hydrogenphosphate (47.0 g) was heated to 45°C in methanol (420 ml). To this mixture sodium methoxide (29 ml) was added and the solution was cooled to room temperature. Separate solutions of 37% aqueous formaldehyde (53 ml) in methanol (7 ml) and sodium borohydride (10.0 g) in water (49 ml stabilised with two drops of 46/48% w/w sodium hydroxide) were then added simultaneously over one hour at a temperature between 17-21°C. Stirring the mixture at room temperature for a further 1 hour was followed by adjustment of the solution to pH 6 by addition of conc. HCl (42 ml) in water (83 ml). The suspension was then removed by filtration and the filter bed washed with methanol (59 ml). Further water (90 ml) was added to the filtrate and then the mixture was

concentrated to remove the residual methanol. Adjustment of the aqueous solution to pH 2 using conc. HCl, addition of ethyl acetate (100 ml) and methanol (10 ml), was followed by basification with potassium carbonate (90 g) in water (130 ml). This precipitated a grey solid that was collected by filtration, washed with water (2 X 100 ml) and then washed with ethyl acetate (100 ml) to give sumatriptan free base (42.4 g, 87%, HPLC showed 98% compound).


Comparative Example.

3-(2-Aminoethyl)-*N*-methyl-1*H*-indole-5-methanesulphonamide was reacted in a similar fashion to example 3 without the inclusion of the sodium hydrogenphosphate buffer. The product free base was obtained in 40% yield and in a purity of 60% by HPLC.


- 1) A process for preparing a compound of formula I including salts thereof



wherein R represents $\text{CH}_2\text{SO}_2\text{NHCH}_3$, $\text{CH}_2\text{CH}_2\text{SO}_2\text{Ph}$, $\text{CH}_2\text{CH}_2\text{SO}_2\text{NHMe}$ or a group of structure:




comprising the reduction of a diazonium salt of formula II




in which R is as previously defined and X^- represents an anion derived from hydrochloric acid, sulphuric acid, acetic acid, phosphoric acid, tetrafluoroboric acid or hydrobromic acid with a dithionite salt.

- 2) A process according to claim 1 wherein the compound of formula II is prepared by reacting a compound of formula III or salt thereof




in which R and X^- represents a chloride, bromide, acetate, hydrosulphate or phosphate anion as previously defined with a diazotising agent optionally in the presence of acid.

- 3) A process according to either claim 1 or claim 2 wherein the compound of formula III is prepared by reacting a compound of formula IV




in which R is as previously defined with a reducing agent, optionally in the presence of an acid.


- 4) A process according to claim 1 wherein the compound of formula I is further reacted with an aldehyde of formula VI in which R1 is as previously defined optionally in the presence of an acid catalyst



to give a compound of formula V



in which R is as previously defined and R1 is a group of formula a), b), c), d), or e):




comprising performing a Fischer Indole synthesis by reacting a compound of formula I with an aldehyde of formula VI in which R1 is as previously defined optionally in the presence of an acid catalyst.

5) A process according to claim 4 in which in the compound of formula I R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and in the compound of formula VI R1 is $\text{CH}_2\text{CH}_2\text{NH}_2$ and in the compound of formula V obtained has R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and R1 is $\text{CH}_2\text{CH}_2\text{NH}_2$.


6) A process according to claim 5 wherein the compound of formula V in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and R1 is $\text{CH}_2\text{CH}_2\text{NH}_2$ is reacted with a formaldehyde equivalent and a reducing agent in the presence of a buffer to give a compound of formula V in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and R1 is $\text{CH}_2\text{CH}_2\text{NMe}_2$.

7) A process for the preparation of sumatriptan or a pharmaceutically acceptable salt thereof comprising the following steps:

a) reducing a compound of formula




in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ with a reducing agent, optionally in the presence of an acid to give a compound of formula III




or optionally a salt thereof in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and

- b) reacting the compound of formula III with a diazotising agent optionally in the presence of acid to give a compound of formula II




in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and X^- represents a chloride, bromide, acetate, hydro-sulphate or phosphate anion and

- c) reducing the compound of formula II with a dithionite salt to give a compound of formula I




in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and

- d) reacting the compound of formula I in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ with an aldehyde of formula VII



VII

in which R_2 represents $\text{ClCH}_2\text{CH}_2-$ and the aldehyde group is protected as an acetal in the presence of a buffer optionally in the presence of an acid catalyst to give a compound of formula V



or a salt thereof in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and R_1 is $\text{CH}_2\text{CH}_2\text{NH}_2$ and

- e) reacting the compound of formula V obtained in d) with a formaldehyde equivalent and a reducing agent in the presence of a buffer to give a compound of formula V in which R is $\text{CH}_2\text{SO}_2\text{NHCH}_3$ and R_1 is $\text{CH}_2\text{CH}_2\text{NMe}_2$ or a pharmaceutically acceptable salt thereof.

- 8) A process according to claim 7 in which none of the intermediate compounds is isolated.
- 9) A process according to claim 7 wherein the salt is the succinate salt.
- 10) A process according to claim 7 wherein the salt is the hemisulphate salt.

INTERNATIONAL SEARCH REPORT

Int lational Application No

PCT/EP 00/10581

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C303/40 C07C311/35 C07D209/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BEILSTEIN Data, WPI Data, EPO-Internal, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

Y	US 4 816 470 A (I.H. COATES, ET AL.) 28 March 1989 (1989-03-28) cited in the application examples 1,21	1-10
---	---	------

Y	US 4 994 483 A (OXFORD ALEXANDER W ET AL) 19 February 1991 (1991-02-19) column 14, line 55 -column 15, line 22; example 1	1-4
---	--	-----

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

8 February 2001

Date of mailing of the international search report

23/02/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

English, R

INTERNATIONAL SEARCH REPORT

Int. :ional Application No

PCT/EP 00/10581

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	L.J. STREET, ET AL.: "Synthesis and serotonergic activity of N,N-dimethyl-2-'5-(1,2,4-triazol-1-ylmethyl)-1H-indol-3-yl!ethylamine and analogues: potent agonists for 5-HT1D reseptors" JOURNAL OF MEDICINAL CHEMISTRY, vol. 38, no. 10, 12 May 1995 (1995-05-12), pages 1799-1810, XP000608327 American Chemical Society, Washington, DC, US ISSN: 0022-2623 scheme 2	1-4
Y	--- R.C. GLENN, ET AL.: "Computer-aided design and synthesis of 5-substituted tryptamines and their pharmacology at the 5-HT1D receptor: discovery of compounds with potential anti-migraine properties" JOURNAL OF MEDICINAL CHEMISTRY, vol. 38, no. 18, 1 September 1995 (1995-09-01), pages 3566-3580, XP002159863 American Chemical Society, Washington, DC, US ISSN: 0022-2623 scheme 5	1-4
Y	--- L. THOMPSON: "The preparation of some hydrazines" JOURNAL OF THE SOCIETY OF DYERS AND COLOURISTS., vol. 37, 1921, pages 7-11, XP000982025 Society of Dyers and Colourists, Bradford, GB ISSN: 0037-9859 the whole document	1-10
A	--- US 5 272 145 A (J.H. HUTCHINSON, ET AL.) 21 December 1993 (1993-12-21) example 1A	1-4
A	--- US 4 851 406 A (A. MERTENS, ET AL.) 25 July 1989 (1989-07-25) examples 1,2	1-4
A	--- GB 2 162 522 A (GLAXO) 5 February 1986 (1986-02-05) cited in the application examples 1-3	6,9,10
A	--- EP 0 490 689 A (GLAXO) 17 June 1992 (1992-06-17) cited in the application the whole document -----	10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/10581

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4816470 A	28-03-1989	AT 381933 B	10-12-1986
		AT 208483 A	15-05-1986
		AU 566149 B	08-10-1987
		AU 1544183 A	15-12-1983
		BE 896986 A	07-12-1983
		CA 1199643 A	21-01-1986
		CH 657359 A	29-08-1986
		CZ 280529 B	14-02-1996
		CY 1458 A	21-07-1989
		DE 3320521 A	08-12-1983
		ES 523039 D	01-10-1984
		ES 8407473 A	16-12-1984
		ES 529802 D	16-11-1985
		ES 8601880 A	01-03-1986
		ES 544281 D	16-07-1986
		ES 8609247 A	16-12-1986
		FI 832035 A,B,	08-12-1983
		FR 2530625 A	27-01-1984
		GB 2124210 A,B	15-02-1984
		GR 79215 A	22-10-1984
		HK 14789 A	24-02-1989
		IE 55358 B	29-08-1990
		IL 68913 A	31-08-1987
		IT 1170387 B	03-06-1987
		JP 1664600 C	19-05-1992
		JP 3029069 B	23-04-1991
		JP 59042366 A	08-03-1984
		KE 3833 A	18-11-1988
		KR 9006681 B	17-09-1990
		LU 84852 A	29-03-1985
		MX 9202623 A	30-06-1992
		MY 17987 A	31-12-1987
		NL 8302031 A,B,	02-01-1984
		NZ 204482 A	10-09-1986
		PH 19908 A	13-08-1986
		PT 76830 A,B	01-07-1983
		SE 452459 B	30-11-1987
		SE 8303208 A	08-12-1983
		SK 404091 A	07-06-1995
		ZA 8304110 A	30-01-1985
US 4994483 A	19-02-1991	AT 381934 B	10-12-1986
		AT 387384 A	15-05-1986
		AU 575365 B	28-07-1988
		AU 3636784 A	13-06-1985
		BE 901224 A	06-06-1985
		CA 1233183 A	23-02-1988
		CH 663411 A	15-12-1987
		CY 1562 A	17-05-1991
		DE 3444572 A	20-06-1985
		DK 214090 A	06-09-1990
		DK 583684 A	07-06-1985
		ES 538336 D	01-06-1986
		ES 8607234 A	01-11-1986
		ES 546631 D	16-10-1987
		ES 8800150 A	01-01-1988
		FI 844789 A,B,	07-06-1985
		FR 2555987 A	07-06-1985

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/10581

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4994483 A		GB 2150932 A,B	10-07-1985
		HK 85690 A	25-10-1990
		HU 40624 A	28-01-1987
		HU 36455 A	30-09-1985
		IL 73756 A	29-02-1988
		IT 1178268 B	09-09-1987
		JP 1878337 C	07-10-1994
		JP 6002733 B	12-01-1994
		JP 60155156 A	15-08-1985
		JP 3184958 A	12-08-1991
		NL 8403719 A	01-07-1985
		NO 844879 A,B,	07-06-1985
		NZ 210469 A	27-01-1989
		PT 80572 A,B	01-06-1985
		SE 458446 B	03-04-1989
		SE 8406200 A	07-06-1985
		SG 65790 G	26-10-1990
		ZA 8409498 A	24-09-1986
US 5272145 A	21-12-1993	US 5204344 A	20-04-1993
		US 5252585 A	12-10-1993
		AU 5620894 A	04-07-1994
		WO 9413293 A	23-06-1994
		US 5380850 A	10-01-1995
		AT 121085 T	15-04-1995
		AU 650185 B	09-06-1994
		AU 3012692 A	11-02-1993
		AU 628212 B	10-09-1992
		AU 6121190 A	28-02-1991
		CA 2023340 A	23-02-1991
		DE 69018553 D	18-05-1995
		DE 69018553 T	19-10-1995
		EP 0419049 A	27-03-1991
		IE 903022 A	27-02-1991
		IL 95371 A	26-08-1994
		JP 2065919 C	24-06-1996
		JP 3163075 A	15-07-1991
		JP 7086101 B	20-09-1995
		NO 176606 B	23-01-1995
		NZ 234883 A	27-01-1995
		NZ 248177 A	27-01-1995
		PT 95055 A	18-04-1991
		ZA 9006610 A	29-05-1991
		CA 2060557 A	06-08-1992
		MX 9203486 A	01-08-1992
		AU 4413893 A	24-01-1994
		WO 9400446 A	06-01-1994
US 4851406 A	25-07-1989	DE 3531658 A	12-03-1987
		AT 73797 T	15-04-1992
		AU 572405 B	05-05-1988
		AU 6216686 A	12-03-1987
		DD 258229 A	13-07-1988
		DE 3684415 A	23-04-1992
		DK 419086 A	06-03-1987
		EP 0223937 A	03-06-1987
		ES 2001936 A	01-07-1988
		FI 863564 A	06-03-1987

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/10581

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4851406 A		GR 862248 A	31-12-1986
		HU 41770 A,B	28-05-1987
		IL 79911 A	15-04-1991
		JP 62056486 A	12-03-1987
		NZ 217419 A	29-03-1989
		PT 83310 A,B	01-10-1986
		ZA 8606705 A	29-04-1987
GB 2162522 A	05-02-1986	CY 1475 A	21-07-1989
		AT 386196 B	11-07-1988
		AT 226685 A	15-12-1987
		AU 573878 B	23-06-1988
		AU 4568985 A	06-02-1986
		BE 903006 A	03-02-1986
		CA 1241004 A	23-08-1988
		CH 666026 A	30-06-1988
		CZ 280530 B	14-02-1996
		DE 3527648 A	13-02-1986
		DK 351185 A,B,	02-02-1986
		EG 17283 A	30-06-1992
		ES 545810 D	16-10-1986
		ES 8700234 A	01-01-1987
		ES 552047 D	16-12-1987
		ES 8801204 A	01-03-1988
		ES 557480 D	16-02-1988
		ES 8801788 A	01-05-1988
		ES 557481 D	16-02-1988
		ES 8801789 A	01-05-1988
		ES 557482 D	01-03-1988
		ES 8801900 A	16-05-1988
		ES 557483 D	16-02-1988
		ES 8801790 A	01-05-1988
		FI 852969 A,B,	02-02-1986
		FR 2568571 A	07-02-1986
		GR 851850 A	26-11-1985
		HK 33289 A	28-04-1989
		HU 40077 A,B	28-11-1986
		IE 58122 B	14-07-1993
		IL 75986 A	28-02-1989
		IT 1207325 B	17-05-1989
		JP 1894608 C	26-12-1994
		JP 6023197 B	30-03-1994
		JP 61047464 A	07-03-1986
		KE 3858 A	02-06-1989
		KR 9303759 B	10-05-1993
		LU 86032 A	18-02-1986
		LU 88266 A	03-02-1994
		MX 159672 A	02-08-1989
		MX 14221 A	01-10-1993
		NL 8502171 A,C	03-03-1986
		NO 853046 A,B,	03-02-1986
		NZ 212948 A	29-05-1989
		PH 22850 A	19-01-1989
		PH 24117 A	05-03-1990
		PL 254800 A	21-10-1986
		PT 80900 B	30-12-1987
		PT 80900 A	01-09-1985
		SE 452460 B	30-11-1987

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/10581

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0490689 A	17-06-1992	AP 230 A	27-01-1993
		AT 119881 T	15-04-1995
		AU 650706 B	30-06-1994
		AU 9072091 A	08-07-1992
		BE 1005085 A	13-04-1993
		CA 2098302 A	13-06-1992
		CH 684192 A	29-07-1994
		CN 1063099 A, B	29-07-1992
		CN 1128259 A, B	07-08-1996
		CZ 9301140 A	16-02-1994
		CY 2005 A	05-12-1997
		DE 69108200 D	20-04-1995
		DE 69108200 T	20-07-1995
		DK 490689 T	22-05-1995
		EG 20219 A	30-11-1997
		WO 9210477 A	25-06-1992
		ES 2069836 T	16-05-1995
		FR 2670487 A	19-06-1992
		GB 2251614 A, B	15-07-1992
		GR 3015430 T	30-06-1995
		HK 82697 A	27-06-1997
		HU 9500422 A	28-09-1995
		IE 914317 A, B	17-06-1992
		IL 100330 A	18-06-1996
		IT 1252868 B	28-06-1995
		JP 2994037 B	27-12-1999
		JP 6503560 T	21-04-1994
		KR 211479 B	02-08-1999
		MX 9102501 A	01-09-1992
		NL 9102071 A	01-07-1992
		NZ 240942 A	25-03-1994
		PT 99757 A, B	31-12-1992
		RU 2108328 C	10-04-1998
		US 5554639 A	10-09-1996
		US 5705520 A	06-01-1998
		ZA 9109750 A	28-10-1992

THIS PAGE BLANK (USPTO)